

Centre Number						Candidate Number			
Surname									
Other Names									
Candidate Signature									

For Examiner's Use	
Examiner's Initials	
Question	Mark
1	
2	
3	
4	
5	
6	
7	
8	
9	
10	
11	
TOTAL	



General Certificate of Education  
Advanced Level Examination  
June 2015

## Chemistry

## CHEM4

### Unit 4 Kinetics, Equilibria and Organic Chemistry

Wednesday 10 June 2015 1.30 pm to 3.15 pm

**For this paper you must have:**

- the Periodic Table/Data Sheet, provided as an insert (enclosed)
- a ruler with millimetre measurements
- a calculator.

**Time allowed**

- 1 hour 45 minutes

**Instructions**

- Use black ink or black ball-point pen.
- Fill in the boxes at the top of this page.
- Answer **all** questions.
- You must answer the questions in the spaces provided. Do not write outside the box around each page or on blank pages.
- All working must be shown.
- Do all rough work in this book. Cross through any work you do not want to be marked.

**Information**

- The marks for questions are shown in brackets.
- The maximum mark for this paper is 100.
- You are expected to use a calculator, where appropriate.
- The Periodic Table/Data Sheet is provided as an insert.
- Your answers to the questions in **Section B** should be written in continuous prose, where appropriate.
- You will be marked on your ability to:
  - use good English
  - organise information clearly
  - use scientific terminology accurately.

**Advice**

- You are advised to spend about 70 minutes on **Section A** and about 35 minutes on **Section B**.



J U N 1 5 C H E M 4 0 1


WMP/Jun15/CHEM4/E5

**CHEM4**

### Section A

Answer **all** questions in the spaces provided.

- 1** Gases **A** and **B** react as shown in the following equation.



The initial rate of the reaction was measured in a series of experiments at a constant temperature. The following rate equation was determined.

$$\text{rate} = k[\mathbf{A}]^2$$

An incomplete table of data for the reaction between **A** and **B** is shown in **Table 1**.

**Table 1**

Experiment	Initial [A] / mol dm <sup>-3</sup>	Initial [B] / mol dm <sup>-3</sup>	Initial rate / mol dm <sup>-3</sup> s <sup>-1</sup>
<b>1</b>	$4.2 \times 10^{-3}$	$2.8 \times 10^{-3}$	$3.3 \times 10^{-5}$
<b>2</b>	$7.9 \times 10^{-3}$	$2.8 \times 10^{-3}$	
<b>3</b>		$5.6 \times 10^{-3}$	$1.8 \times 10^{-4}$

- 1 (a)** Use the data from Experiment **1** to calculate a value for the rate constant, *k*, at this temperature.  
 Deduce the units of *k*.

**[3 marks]**

Calculation .....

.....

.....

.....

Units .....

.....



- 1 (b) Use your value of  $k$  from Question 1(a) to complete **Table 1** for the reaction between **A** and **B**.

(If you have been unable to calculate an answer for Question 1 (a), you may assume a value of 2.3. This is **not** the correct answer.)

[2 marks]

- 1 (c) The reaction is zero order with respect to **B**.

State the significance of this zero order for the mechanism of the reaction.

[1 mark]

.....  
.....  
.....

6


Turn over for the next question

Turn over ►



0 3

- 2 Sulfur dioxide reacts with oxygen to form sulfur trioxide according to the equation



- 2 (a) Write an expression for the equilibrium constant,  $K_c$ , for this reaction and deduce its units.

[2 marks]

$K_c$  .....

.....

Units .....

.....

- 2 (b) Samples of sulfur dioxide, oxygen and sulfur trioxide were added to a flask of volume 1.40 dm<sup>3</sup> and allowed to reach equilibrium at a given temperature.

The flask contained 0.0550 mol of sulfur dioxide and 0.0720 mol of sulfur trioxide at equilibrium.

$K_c$  has the numerical value of 27.9 under these conditions.

Calculate the amount, in moles, of oxygen gas in this equilibrium mixture.

[3 marks]

.....

.....

.....

.....

.....

.....

.....

.....



**2 (c)** The experiment in Question **2 (b)** was repeated with the same amounts of sulfur dioxide, oxygen and sulfur trioxide at the same temperature but in a smaller flask. The mixture was allowed to reach equilibrium.

**2 (c) (i)** State the effect, if any, of using a smaller flask on the value of  $K_c$

[1 mark]

.....

**2 (c) (ii)** State the effect, if any, of using a smaller flask on the amount of sulfur trioxide at equilibrium.

Explain your answer.

[3 marks]

Effect .....

Explanation .....

.....

.....

.....

9

Turn over for the next question

Turn over ►




0 5

WMP/Jun15/CHEM4

- 3 Titration curves, labelled **E**, **F**, **G** and **H**, for combinations of different aqueous solutions of acids and bases are shown in **Figure 1**.

All solutions have concentrations of  $0.1 \text{ mol dm}^{-3}$ .

**Figure 1**



- 3 (a) In this part of the question, write the appropriate letter in each box.

From the curves **E**, **F**, **G** and **H**, choose the curve produced by the addition of

- 3 (a) (i) sodium hydroxide to  $25 \text{ cm}^3$  of ethanoic acid

[1 mark]

- 3 (a) (ii) ammonia to  $25 \text{ cm}^3$  hydrobromic acid

[1 mark]

- 3 (a) (iii) hydrochloric acid to  $25 \text{ cm}^3$  of potassium hydroxide

[1 mark]



0 6

- 3 (b) **Table 2** shows information about some acid-base indicators.

**Table 2**

Indicator	pH range	Lower pH colour	Higher pH colour
pentamethoxy red	1.2–3.2	violet	colourless
naphthyl red	3.7–5.0	red	yellow
4-nitrophenol	5.6–7.0	colourless	yellow
cresol purple	7.6–9.2	yellow	purple

- 3 (b) (i) Which indicator in **Table 2** could be used for the titration that produces curve **E** but not for the titration that produces curve **F**?

[1 mark]

Tick ( $\checkmark$ ) one box.

pentamethoxy red

naphthyl red

4-nitrophenol

cresol purple

- 3 (b) (ii) Give the colour change at the end point of the titration that produces curve **H** when naphthyl red is used as the indicator.

[1 mark]

.....

- 3 (b) (iii) A beaker contains 25 cm<sup>3</sup> of a buffer solution at pH = 6.0. Two drops of each of the four indicators in **Table 2** are added to this solution.

State the colour of the mixture of indicators in this buffer solution.  
You should assume that the indicators do **not** react with each other.

[1 mark]

.....


6

Turn over ►



0 7

- 4 Water dissociates slightly according to the equation:



The ionic product of water,  $K_w$ , is given by the expression

$$K_w = [\text{H}^+][\text{OH}^-]$$

$K_w$  varies with temperature as shown in **Table 3**.

**Table 3**

Temperature / °C	$K_w$ / mol <sup>2</sup> dm <sup>-6</sup>
25	$1.00 \times 10^{-14}$
50	$5.48 \times 10^{-14}$

- 4 (a) Explain why the expression for  $K_w$  does **not** include the concentration of water. [2 marks]

.....  
.....  
.....  
.....  
.....  
.....

- 4 (b) Explain why the value of  $K_w$  increases as the temperature increases. [2 marks]

.....  
.....  
.....  
.....  
.....



- 4 (c) Calculate the pH of pure water at 50 °C.  
Give your answer to 2 decimal places.

[3 marks]

.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....

- 4 (d) Calculate the pH of 0.12 mol dm<sup>-3</sup> aqueous NaOH at 50 °C.  
Give your answer to 2 decimal places.

[3 marks]

.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....

10


Turn over for the next question

Turn over ►



0 9

- 5 Consider the following reaction sequence starting from methylbenzene.



J

- 5 (a) Name the type of mechanism for reaction 1.

[1 mark]

---

- 5 (b) Compound J is formed by reduction in reaction 2.

- 5 (b) (i) Give a reducing agent for this reaction.

[1 mark]

---

- 5 (b) (ii) Write an equation for this reaction. Use [H] to represent the reducing agent.

[1 mark]

---

- 5 (b) (iii) Give a use for J.

[1 mark]

---



1 0

- 5 (c) Outline a mechanism for the reaction of bromomethane with an excess of compound **J**.  
You should represent **J** as  $\text{RNH}_2$  in the mechanism.

[4 marks]

- 5 (d) Compound **K** ( $\text{C}_6\text{H}_5\text{CH}_2\text{NH}_2$ ) is a structural isomer of **J**.

Explain why **J** is a weaker base than **K**.

[3 marks]

.....

.....

.....

.....

.....

.....



Turn over ►



**6** Esters are used as raw materials in the production of soaps and biodiesel.

**6 (a)** A student prepared an ester by two different methods.

Method 1 alcohol + acid anhydride

Method 2 alcohol + acyl chloride

**6 (a) (i)** An ester was prepared using method 1, by reacting  $(CH_3)_2CHOH$  with  $(CH_3CO)_2O$

Write an equation for this reaction and give the IUPAC name of the ester formed.

**[2 marks]**

Equation

IUPAC name of the ester .....

**6 (a) (ii)** The same ester was prepared using method 2 by reacting  $(CH_3)_2CHOH$  with  $CH_3COCl$

Outline a mechanism for this reaction.


**[4 marks]**



1 2


- 6 (b)** The ester shown occurs in vegetable oils.

It can be hydrolysed to make soap and can also be used to produce biodiesel.



- 6 (b) (i)** Write an equation for the reaction of this ester with sodium hydroxide to form soap.

[2 marks]



- 6 (b) (ii)** Give the formula of the biodiesel molecule with the highest  $M_r$  that can be produced by reaction of this ester with methanol.

[1 mark]

.....

9


**Turn over for the next question**

**Turn over ►**




1 3

- 7 (a) The structures and common names of two amino acids are shown.



proline



alanine

- 7 (a) (i) Draw the structure of the zwitterion of proline.

[1 mark]



- 7 (a) (ii) Draw the structure of the tripeptide formed when a proline molecule bonds to two alanine molecules, one on each side.

[2 marks]



1 4

7 (b) Sections of two polymers, **L** and **M**, are shown.



7 (b) (i) Give the IUPAC name of a monomer that forms polymer **L**.

[1 mark]

.....

7 (b) (ii) Give the IUPAC name of the monomer that forms polymer **M**.

[1 mark]

.....

7 (b) (iii) Draw the section of a polymer made from a dicarboxylic acid and a diamine that is isomeric with the section of polymer **M** shown.

[1 mark]

.....

7 (b) (iv) Explain why polymer **L** is non-biodegradable.

[1 mark]



.....

.....

.....



- 8** Equations for the hydrogenation of cyclohexene and of benzene, together with the enthalpies of hydrogenation, are shown.



- 8 (a) (i)** Use these data to show that benzene is  $152 \text{ kJ mol}^{-1}$  more stable than the hypothetical compound cyclohexa-1,3,5-triene.

[1 mark]


.....  
.....  
.....  
.....

- 8 (a) (ii)** State, in terms of its bonding, why benzene is more stable than cyclohexa-1,3,5-triene.

[1 mark]

.....  
.....  
.....  
.....

- 8 (b)** Three carbon-carbon bonds are labelled on the structures shown. These bonds are of different lengths.




Write the letters **w**, **x** and **y** in order of **increasing** bond length.

[1 mark]


.....



- 8 (c)** The structures of two cyclic dienes are shown.



cyclohexa-1,4-diene



cyclohexa-1,3-diene

- 8 (c) (i)** Use the enthalpy of hydrogenation data given opposite to calculate a value for the enthalpy of hydrogenation of cyclohexa-1,4-diene.

[1 mark]

.....  
.....

- 8 (c) (ii)** Predict a value for the enthalpy of hydrogenation of cyclohexa-1,3-diene.

[1 mark]

.....

- 8 (c) (iii)** Explain your answers to Questions **8 (c) (i)** and **8 (c) (ii)** in terms of the bonding in these two dienes.

[3 marks]

.....  
.....  
.....  
.....  
.....  
.....

8

Turn over ►



**Section B**

Answer **all** questions in the spaces provided.

**9**

The acid dissociation constant,  $K_a$ , for ethanoic acid is given by the expression

$$K_a = \frac{[H^+][\text{CH}_3\text{COO}^-]}{[\text{CH}_3\text{COOH}]}$$

The value of  $K_a$  for ethanoic acid is  $1.74 \times 10^{-5}$  mol dm $^{-3}$  at 25 °C.

**9 (a)**

A buffer solution is prepared using ethanoic acid and sodium ethanoate. In the buffer solution, the concentration of ethanoic acid is 0.186 mol dm $^{-3}$  and the concentration of sodium ethanoate is 0.105 mol dm $^{-3}$ .

Calculate the pH of this buffer solution.

Give your answer to 2 decimal places.

**[3 marks]**

.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....



- 9 (b)** In a different buffer solution, the concentration of ethanoic acid is  $0.251 \text{ mol dm}^{-3}$  and the concentration of sodium ethanoate is  $0.140 \text{ mol dm}^{-3}$ .

A sample of hydrochloric acid containing  $0.015 \text{ mol}$  of HCl is added to  $1000 \text{ cm}^3$  of this buffer solution.

Calculate the pH of the buffer solution after the hydrochloric acid has been added.  
You should ignore any change in total volume.  
Give your answer to 2 decimal places.

[5 marks]

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---






8

**Turn over for the next question**

**Turn over ►**



- 10** The following five isomers, **P**, **Q**, **R**, **S** and **T**, were investigated using test-tube reactions and also using n.m.r. spectroscopy.

**P****Q****R****S****T**

- 10 (a)** A simple test-tube reaction can be used to distinguish between isomers **P** and **S**.

Identify a reagent (or combination of reagents) you could use.

State what you would observe when both isomers are tested separately with this reagent or combination of reagents.

[3 marks]

---



---



---



---



---



- 10 (b) A simple test-tube reaction can be used to distinguish between isomer **Q** and all the other isomers.

Identify a reagent (or combination of reagents) you could use.

State what you would observe when **Q** is tested with this reagent or combination of reagents.

[2 marks]

.....  
.....  
.....  
.....

- 10 (c) State which **one** of the isomers, **P**, **Q**, **R**, **S** and **T**, has the least number of peaks in its  $^1\text{H}$  n.m.r. spectrum.

Give the number of peaks for this isomer.

[2 marks]

.....  
.....  
.....

- 10 (d) Write the **molecular** formula of the standard used in  $^{13}\text{C}$  n.m.r. spectroscopy.  
Give **two** reasons why this compound is used.

[3 marks]



.....  
.....  
.....  
.....  
.....  
.....

Question 10 continues on the next page






Turn over ►



10 (e) **Figure 2** and **Figure 3** show the  $^{13}\text{C}$  n.m.r. spectra of two of the five isomers.

**Figure 2****Figure 3**

The structures of the five isomers are repeated to help you answer this question.

**P****Q****R****S****T**

2 2

State which isomer produces the spectrum in **Figure 2** and which isomer produces the spectrum in **Figure 3**.

Explain your answer.

You do not need to identify every peak in each spectrum.  
Use **Table C** on the Data Sheet to answer the question.

[5 marks]

.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....

- 10 (f)** **U** and **V** are other isomers of **P**, **Q**, **R**, **S** and **T**.  
The  $^1\text{H}$  n.m.r. spectrum of **U** consists of two singlets.  
**V** is a cyclic alcohol that exists as optical isomers.

Draw the structure of **U** and the structure of **V**.

[2 marks]

**U**

**V**

17

Turn over ►



2 3

**11**

The N-substituted amide  $C_6H_{13}NO$  can be formed from but-2-ene in a three-step synthesis.



For each reaction

- state a reagent
- give the structure of the product
- name the mechanism of the reaction.

Detailed mechanisms are **not** required.

**[9 marks]**

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

**END OF QUESTIONS**

**9**

