

# Mark Scheme (Results) January 2011

GCE

GCE Chemistry (6CH04/01)

Edexcel is one of the leading examining and awarding bodies in the UK and throughout the world. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers.

Through a network of UK and overseas offices, Edexcel's centres receive the support they need to help them deliver their education and training programmes to learners.

For further information, please call our GCE line on 0844 576 0025, our GCSE team on 0844 576 0027, or visit our website at [www.edexcel.com](http://www.edexcel.com).

If you have any subject specific questions about the content of this Mark Scheme that require the help of a subject specialist, you may find our Ask The Expert email service helpful.

Ask The Expert can be accessed online at the following link:

<http://www.edexcel.com/Aboutus/contact-us/>

Alternatively, you can speak directly to a subject specialist at Edexcel on our dedicated Science telephone line: 0844 576 0037

January 2011

Publications Code UA026200

All the material in this publication is copyright  
© Edexcel Ltd 2011

### Section A (multiple choice)

Question Number	Correct Answer	Mark
1 (a)	D	1

Question Number	Correct Answer	Mark
1 (b)	B	1

Question Number	Correct Answer	Mark
1 (c)	A	1

Question Number	Correct Answer	Mark
2	D	1

Question Number	Correct Answer	Mark
3	D	1

Question Number	Correct Answer	Mark
4	B	1

Question Number	Correct Answer	Mark
5	C	1

Question Number	Correct Answer	Mark
6	C	1

Question Number	Correct Answer	Mark
7 (a)	A	1

Question Number	Correct Answer	Mark
7 (b)	A	1

Question Number	Correct Answer	Mark
7 (c)	D	1

Question Number	Correct Answer	Mark
8	B	1

Question Number	Correct Answer	Mark
9	D	1

Question Number	Correct Answer	Mark
10	C	1

Question Number	Correct Answer	Mark
11	D	1

Question Number	Correct Answer	Mark
12	D	1

Question Number	Correct Answer	Mark
13	A	1

Question Number	Correct Answer	Mark
14	B	1

Question Number	Correct Answer	Mark
15	C	1

Question Number	Correct Answer	Mark
16	A	1

**TOTAL FOR SECTION A = 20 MARKS**

Section B

Question Number	Acceptable Answers	Reject	Mark
17 (a)	$\Delta S_{\text{system}} = (3 \times 2 \times 65.3 + 197.6) - (186.2 + 188.7)$ Correct data for CH <sub>4</sub> and CO (186.2 and 197.6) (1) = (+) 214.5 / 215 (J mol <sup>-1</sup> K <sup>-1</sup> ) / (+) 0.2145 / 0.215 kJ (mol <sup>-1</sup> K <sup>-1</sup> ) (1) Units must be shown if data has been converted to kJ Full marks (2) for correct answer without working Ignore sf except 1 Answer of -214.5 scores (1) Answer of +18.6 if entropy of H not doubled scores (1) Answer of -46.7 if entropy of H <sub>2</sub> not tripled scores (1) ALLOW TE in second mark for minor error in data e.g. writing 63.5 instead of 65.3. No TE if data used is not entropy of compounds.	214 0.214	2

Question Number	Acceptable Answers	Reject	Mark
17 (b)	$(\Delta S_{\text{surroundings}}) = \frac{-\Delta H}{T}$ Expression or use of expression, $\frac{-206.1 \times (1000)}{298}$ (1) = -691.6 J (mol <sup>-1</sup> K <sup>-1</sup> ) / -0.6916 kJ (mol <sup>-1</sup> K <sup>-1</sup> ) (1) Ignore sf except 1		2

Question Number	Acceptable Answers	Reject	Mark
17 (c)	$\Delta S_{\text{total}} = (214.5 + (-691.6)) = -477.1 \text{ (J mol}^{-1} \text{ K}^{-1}) /$ $- 0.4771 \text{ (kJ mol}^{-1} \text{ K}^{-1}) \text{ (1)}$ <p>ALLOW TE for answer to (a) plus answer to (b). If 214.5 is added to -0.69 no TE unless -0.69 is specified to be in joules. Ignore sf except 1</p> <p>Negative / less than zero (so not spontaneous) / would be positive if spontaneous. (1)</p> <p>ALLOW "feasible" for spontaneous.</p> <p>If answer to calculation is positive, accept comment that it would be expected to be negative if not spontaneous</p>	<p>Addition of value in J to specified value in kJ</p> <p>Comments on kinetic stability</p>	2

Question Number	Acceptable Answers	Reject	Mark										
*17 (d) (i)	<p><math>K_p = \frac{(p_{H_2})^3 \times (p_{CO})}{(p_{CH_4})(p_{H_2O})}</math> (1)</p> <p>4 Correct partial pressures (3)</p> <table border="1" data-bbox="296 432 761 607"> <tr> <td></td> <td>CH<sub>4</sub></td> <td>H<sub>2</sub>O</td> <td>CO</td> <td>H<sub>2</sub></td> </tr> <tr> <td>pp</td> <td>0.25</td> <td>0.25</td> <td>0.375</td> <td>1.125</td> </tr> </table> <p>ALLOW partial pressures as fractions</p> <p><math>K_p = \frac{(1.125)^3 \times (0.375)}{(0.25)(0.25)} = 8.54 \text{ atm}^2</math></p> <p>value of <math>K_p</math> (1)</p> <p>unit (1) (Stand alone mark)</p> <p>Correct calculation without working scores the 5 calculation marks.</p> <p>TE from <math>K_p</math> expression if inverted Ignore sf except 1</p> <p><b>If any partial pressures are incorrect:</b> Calculating total number of moles (6.4) (1)</p> <p>Calculating mole fractions (0.125, 0.125, 0.1875, 0.5625 if total number of moles is correct) (1)</p> <p>Multiplying mole fractions by total pressure (x 2 atm) (1)</p> <p>value of <math>K_p</math> (1)</p> <p>unit (1) (stand alone mark)</p> <p>ALLOW TE in value of <math>K_p</math> only from incorrect partial pressures, not using values in question as not using equilibrium moles</p> <p>If treated as a <math>K_c</math> calculation following <math>K_p</math> expression : <math>K_p</math> expression (1) units <math>\text{atm}^2</math> (1)</p> <p>Max. mark (2)</p>		CH <sub>4</sub>	H <sub>2</sub> O	CO	H <sub>2</sub>	pp	0.25	0.25	0.375	1.125	<p>Square brackets</p> <p>TE for <math>K_p</math> expression with addition, not multiplication</p>	<p>6</p>
	CH <sub>4</sub>	H <sub>2</sub> O	CO	H <sub>2</sub>									
pp	0.25	0.25	0.375	1.125									

Question Number	Acceptable Answers	Reject	Mark
17 (d) (ii)	$\Delta S_{\text{total}} = (8.31 \ln 8.54) = (+)17.8 \text{ (J mol}^{-1} \text{ K}^{-1}\text{)}$ Accept any value that rounds to 17.8  TE from value in (i)  $K_p$ value of 87.48 (obtained by treating calculation in (i) as $K_c$ ) gives $\Delta S_{\text{total}} = 37.16 / 37.12$		1

Question Number	Acceptable Answers	Reject	Mark
17 (d) (iii)	$17.8 = 225 - \frac{206.1 \times 1000}{T} \quad (1)$ $T = \frac{(206.1 \times 1000)}{207.2} = 995 / 990 \text{ (K)} \quad (1)$  Correct answer with no working shown scores 2 Correct method with wrong answer or missing $10^3$ scores 1  TE from (ii) $K_p$ value of 87.48 gives $T = 1097$  <b>OR</b>  If $\Delta S_{\text{total}}$ is taken as zero $0 = 225 - \frac{206.1 \times 1000}{T} \quad (1)$ $T = 916\text{K} \quad (1)$  $K_p$ value of 87.48 gives $T = 916$  Ignore sf except 1		2





Question Number	Acceptable Answers	Reject	Mark
18 (a)	pH = (-log 0.25) = 0.602 / 0.60 / 0.6 Ignore significant figures		1

Question Number	Acceptable Answers	Reject	Mark
18 (b) (i)	$K_a = \frac{[H^+][CH_3CH_2COO^-]}{[CH_3CH_2COOH]}$ <p>ALLOW <math>[H_3O^+]</math> for <math>[H^+]</math></p> <p>ALLOW <math>C_2H_5</math> for <math>CH_3CH_2</math></p> <p>ALLOW <math>\frac{[H^+][A^-]}{[HA]}</math> if HA and A<sup>-</sup> identified</p>	<p>Wrong / missing charge on <math>CH_3CH_2COO^-</math></p> $K_a = \frac{[H^+]^2}{[CH_3CH_2COOH]}$ <p>unless full expression also given</p>	1

Question Number	Acceptable Answers	Reject	Mark
18 (b) (ii)	$1.3 \times 10^{-5} = \frac{[H^+]^2}{0.25}$ / rearrangement of this expression (1) <p><math>([H^+] = 1.8 \times 10^{-3})</math></p> <p>pH = 2.74 (1)</p> <p>Correct answer with no working scores (2) No TE for incorrect <math>[H^+]</math></p> <p>Ignore significant figures except 1 Minimum of 1 decimal place needed</p>		2

Question Number	Acceptable Answers	Reject	Mark
18 (c) (i)	$CH_3CH_2COOH + NaOH \rightarrow CH_3CH_2COO^{(-)}Na^{(+)} + H_2O$ <p>OR <math>CH_3CH_2COOH + OH^- \rightarrow CH_3CH_2COO^- + H_2O</math></p> <p>Accept <math>CH_3CH_2CO_2H</math>, <math>C_2H_5COOH</math>, <math>C_2H_5CO_2H</math></p>	Equations for ethanoic acid	1

Question Number	Acceptable Answers	Reject	Mark
18 (c) (ii)	<p><math>1.3 \times 10^{-5} = \frac{[\text{H}^+][5 \times 10^{-2}]}{[7.5 \times 10^{-2}]}</math> (concentration ratio)</p> <p>OR</p> <p><math>1.3 \times 10^{-5} = \frac{[\text{H}^+](1 \times 10^{-3})}{(1.5 \times 10^{-3})}</math> (ratio by moles)  (ratio by moles allowed as volumes acid and salt equal) (1)</p> <p>( <math>[\text{H}^+] = 1.95 \times 10^{-5}</math> )</p> <p>pH = 4.7 / 4.7099654 (1)</p> <p>Second mark dependent on first  Correct answer with or without working (2)</p> <p>OR</p> <p>pH = pK<sub>a</sub> -log <math>\frac{(1.5 \times 10^{-3})}{1 \times 10^{-3}}</math></p> <p>OR</p> <p>pH = pK<sub>a</sub> -log <math>\frac{(7.5 \times 10^{-2})}{5 \times 10^{-2}}</math> (1)</p> <p>pH = 4.7 (1)</p> <p>Correct answer with or without working (2)</p> <p>Accept any value which rounds to 4.7</p>		2

Question Number	Acceptable Answers	Reject	Mark
*18 (c) (iii)	<p>Mixture is a buffer (1)</p> <p>EITHER</p> <p>OH<sup>-</sup> combines with H<sup>+</sup> in solution (1)</p> <p>Propanoic acid dissociates to replace H<sup>+</sup> (1) <i>Correct equations could gain these marks</i></p> <p>OR</p> <p>OH<sup>-</sup> reacts with propanoic acid (1) <i>Correct equation could gain this mark</i></p> <p>Significant quantities of weak acid and salt are both present /ratio of acid and salt does not change (1)</p> <p>ALLOW a reservoir of weak acid and salt are present: Allow conjugate base for salt</p>	NaOH combines	3

Question Number	Acceptable Answers	Reject	Mark
18 (c) (iv)	<p>S-shaped curve, vertical at 25 cm<sup>3</sup> (with kink at start ) (1)</p> <p>Starting at pH 2-3 (TE from (b)(ii), finishing at pH 12 -13 (1)</p> <p>Vertical section between 3 and 6 units high centred round a pH of between 8 and 9 (1)</p> <p>Vertical section should not extend over more than ±2.5cm<sup>3</sup> This section should start between 5.5 and 7.5 and finish between 9.5 and 11.5 but do not penalise for very small differences.</p> <p>Reverse curve maximum 2</p>		3

Question Number	Acceptable Answers	Reject	Mark
18 (c) (v)	<p>Either Need indicator changing in vertical region of curve / need indicator changing where pH changes sharply / bromocresol green changes before the vertical region (1)</p> <p><b>Not</b> bromocresol green which changes at <b>3.8 - 5.4</b> (1)</p> <p>OR</p> <p><math>pK_{in} \pm 1</math> must be in vertical section / sharply changing section (1)</p> <p><b>Not</b> bromocresol green because <math>pK_{in}</math> is <b>4.7</b> (1)</p> <p>TE from curve with vertical section including pH 3.7 - 5.7</p>	Just “the equivalence point is outside the bromocresol green range”	2

Question Number	Acceptable Answers	Reject	Mark
18 (d) (i)	Dilute acid / dilute strong named acid or formula / NaOH(aq) followed by dilute acid / water plus dilute acid / water plus $H^+$	NaOH alone  water any weak acid concentrated sulfuric acid HCN acid hydrolysis alone	1

Question Number	Acceptable Answers	Reject	Mark
18 (d) (ii)	$CH_3CH_2COCl + H_2O \rightarrow CH_3CH_2COOH + HCl$ / $C_2H_5COCl + H_2O \rightarrow C_2H_5COOH + HCl$  Accept displayed formula	Equations with NaOH or $OH^-$	1

Question Number	Acceptable Answers	Reject	Mark
18 (d) (iii)	Colour change orange to green / blue		1

Question Number	Acceptable Answers	Reject	Mark
18 (e)	<p>Reducing agent /Reduction (of the acid) occurs (1)</p> <p>Li Al H<sub>4</sub> / lithium tetrahydridoaluminate / lithium aluminium hydride (1)</p> <p>Allow minor error in name if correct formula is given</p> <p>Ignore solvent</p> <p>ALLOW nucleophile AND H<sup>-</sup> for 1 mark</p>	<p>Lithal without correct name or formula</p>	2

Question Number	Acceptable Answers	Reject	Mark
19 (a)	<p>Quenches reaction / stops reaction / slows reaction / freezes reaction (1)</p> <p>EITHER by neutralizing the acid / removing the acid / neutralizing the catalyst / removing the catalyst</p> <p>OR</p> <p>So that the acid does not react with the thiosulfate (1)</p>	<p>By neutralizing HI Just “by diluting the reaction mixture” just “by neutralizing the reaction mixture”</p>	2

Question Number	Acceptable Answers	Reject	Mark
19 (b)	Starch (solution)		1

Question Number	Acceptable Answers	Reject	Mark
19 (c)	<p><b>First mark</b> So that [propanone] and [acid] are (virtually) constant</p> <p><b>OR</b> so that the [propanone] and [H<sup>+</sup>] do not affect the rate</p> <p><b>OR</b> Propanone and acid are in excess so <b>changes in concentration</b> don't affect rate (1)</p> <p><b>Second mark</b> And therefore rate changes would only depend on [iodine]</p> <p><b>OR</b> so that the overall order is not determined</p> <p>ALLOW [Iodine] is the limiting factor (1)</p> <p><b>NOTE</b> "so that only the [I<sub>2</sub>] changes" scores (2)</p> <p>"so that only the I<sub>2</sub> concentration changes" scores (2)</p> <p>"so that only the I<sub>2</sub> changes" scores (1)</p>	Propanone and acid are in excess, without reference to further comments	2

Question Number	Acceptable Answers	Reject	Mark
19 (d)	<p>Zero order (1)</p> <p>(Gradient =) <b>rate</b> is constant / I<sub>2</sub> (concentration) doesn't affect <b>rate</b> / <b>rate</b> of change of I<sub>2</sub> (concentration) doesn't change with time (1)</p> <p><b>Mark independently</b></p>	<p>Just 'straight line' Or just 'gradient is constant'</p> <p>[Thiosulfate] or volume of Thiosulfate is proportional to time without reference to iodine</p> <p>Reference to half life [I<sub>2</sub>] is proportional to rate</p>	2



Question Number	Acceptable Answers	Reject	Mark
19 (e)	<p>Measuring cylinder quicker / Measuring cylinder can measure a variety of volumes (1)</p> <p>ALLOW Measuring cylinder can be plastic so unbreakable Comment on lower cost of measuring cylinder if qualified with a reason</p> <p>Pipette more accurate / (graduated) pipette more precise / pipette can be used to extract samples from a reaction mixture (for titration) (1)</p>	<p>Just “Measuring cylinder easier to use” Easier to clean</p> <p>Measuring cylinder can be used for large volumes</p> <p>Pipette more reliable</p> <p>Ignore references to easier</p>	2

Question Number	Acceptable Answers	Reject	Mark
19 (f) (i)	To keep (total) volume constant / to make the (total) volume 32 cm <sup>3</sup> / to make concentrations proportional to volume of reactant	To keep concentrations constant	1

Question Number	Acceptable Answers	Reject	Mark
19 (f) (ii)	<p>First order wrt propanone with explanation (1)</p> <p>First order wrt hydrogen ions/ sulfuric acid, with explanation (1)</p> <p>Explanation can be in terms of experiments 1 and 3 (propanone) or 1 and 2 (acid) and can be in terms of concentration or volume</p> <p>Rate = <math>k[\text{CH}_3\text{COCH}_3][\text{H}^+][\text{I}_2]^0</math> / Rate = <math>k[\text{CH}_3\text{COCH}_3][\text{H}_2\text{SO}_4][\text{I}_2]^0</math> (1)</p> <p>ALLOW names of propanone and sulfuric acid in place of formulae</p> <p>Ignore case of k in rate equation</p> <p>Ignore order wrt iodine even if wrong</p> <p>Third mark is consequential if incorrect orders of propanone and acid given.</p>	<p>Expressions without rate or k</p> <p>Expressions with <math>K_c</math></p> <p>R / r for rate</p>	3

TOTAL FOR SECTION B = 50 MARKS

Section C

Question Number	Acceptable Answers	Reject	Mark
20 (a)	<p>Q: O-H</p> <p>ALLOW OH - O - H (1)</p> <p>R: C=O</p> <p>ALLOW - C = O   - C = O (1)</p> <p>IGNORE names ACCEPT answers written on spectrum</p>	<p>Just 'alcohol' - OH</p> <p>Just 'carbonyl' - C O   C-O</p>	2

Question Number	Acceptable Answers	Reject	Mark
20 (b) (i)	<p>Y = methanol / CH<sub>3</sub>OH (1)</p> <p>Any two of the following: Molecular ion / M<sup>+</sup> / M<sub>r</sub> / CH<sub>3</sub>OH<sup>+</sup> / methanol = 32 CH<sub>3</sub><sup>+</sup> = 15 CH<sub>3</sub>O<sup>+</sup> / CH<sub>2</sub>OH<sup>+</sup> = 31 CHOH<sup>+</sup> / CH<sub>2</sub>O<sup>+</sup> = 30 COH<sup>+</sup> = 29 CO<sup>+</sup> = 28 (1)</p> <p>Charges not required</p> <p>TE in second mark for two correct possible peaks from an incorrect compound.</p>		2

Question Number	Acceptable Answers	Reject	Mark
20 (b) (ii)	<p>Two (1) This mark may be scored if two shifts are given.</p> <p>Any two shifts correctly identified: -OH at 2.0-4.0 / any value in this range H-C-O at 3.0- 4.2 / any value in this range H in CH<sub>3</sub> OH at 3.39 (ppm) (1)</p> <p>Allow TE for ethanol with three peaks (1) and three correct shift values: -OH at 2.0-4.0 / any value in this range H-C-O at 3.0- 4.2 / any value in this range CH in an alkane at 0.1-1.9 (1)</p>	<p>CH in an alkane at 0.1-1.9</p> <p>Just CH<sub>3</sub> OH at 3.39</p>	2

Question Number	Acceptable Answers	Reject	Mark
20 (c) (i)	<p>Z contains two -OH/ one alcohol + one acid</p> <p>ALLOW two alcohol groups / is a diol</p>		1

Question Number	Acceptable Answers	Reject	Mark
20 (c) (ii)	<p>Z is an acid / contains -COOH / contains -CO<sub>2</sub>H/ contains a carboxylic acid group / contains H<sup>+</sup></p>		1

Question Number	Acceptable Answers	Reject	Mark
20 (c) (iii)	<p>Z is a secondary alcohol/ a ketone is <b>formed</b> from Z /</p> <p>Z contains <math>\begin{array}{c}   \\ -\text{C}-\text{OH} \\   \\ \text{H} \end{array}</math> (1)</p>	Z is a ketone	1

Question Number	Acceptable Answers	Reject	Mark
20 (c) (iv)	<p>(Iodoform produced ) so Z contains CH<sub>3</sub> CH(OH)-</p> <p>TE if Z is identified as a ketone in (iii): Z contains CH<sub>3</sub> C=O / Z is a methyl ketone</p>		1

Question Number	Acceptable Answers	Reject	Mark
20 (d)	<p>Answers will be based on several pieces of information (molecular formula, products of ester hydrolysis, answers to (c)) which may be contradictory if errors have been made.</p> <p>ALLOW TE marks for formulae which are chemically possible (ie no 5 bonded carbons etc) and based on most of the deductions but <b>not necessarily all</b>.</p> <p>Z is <math>\text{CH}_3\text{CH}(\text{OH})\text{CH}_2\text{COOH}</math> (1) Stand alone mark</p> <p>ALLOW TE for an acid with OH in wrong position in Z if oxidation product identified as aldehyde</p> <p>TE for Z = <math>\text{CH}_3\text{COCH}_2\text{COOH}</math> if identified as ketone in (iii)</p> <p>X is <math>\text{CH}_3\text{CH}(\text{OH})\text{CH}_2\text{COOCH}_3</math> (1) Stand alone mark TE for a methyl ester of Z</p>		2

Question Number	Acceptable Answers	Reject	Mark
21 (a) (i)	Transesterification Ethanol transesterification	Substituted esterification	1

Question Number	Acceptable Answers	Reject	Mark
21 (a) (ii)	To prevent hydrolysis/ to stop fatty acids forming / to stop breakdown of esters / water reacts with esters/ water is a better nucleophile than ethanol	To dilute ethanol Ethanol would react with water A reaction would occur (unspecified)	1

Question Number	Acceptable Answers	Reject	Mark
21 (b)	<p>(Vegetable) Fats/ oils are renewable (crude oil is not) /</p> <p>biodiesel comes from a renewable source /</p> <p>doesn't use up fossil fuel resources/</p> <p>carbon footprint is less /</p> <p>(closer to) carbon neutral / growing vegetables absorb CO<sub>2</sub></p> <p>If more than one answer is given, and one is incorrect, no mark</p> <p>Ignore comments on biodegradability</p>	<p>Just "made from plants"</p> <p>Just "crude oil is not sustainable"</p> <p>Less polluting produces less greenhouse gases / less CO<sub>2</sub></p> <p>Burns more cleanly</p> <p>Requires less energy for production</p>	1

Question Number	Acceptable Answers	Reject	Mark
21 (c)	<p>Substances to be separated have different (forces of) attraction to / affinity for / solubilities in / adsorption to one or both of the mobile and stationary phases OWTTE (1)</p> <p>ALLOW absorption</p> <p>GC: <b>mobile</b> phase a (inert / unreactive) <b>gas</b> OR GC: <b>mobile</b> phase nitrogen / helium / argon / other named inert gas (1)</p> <p>GC: <b>Stationary</b> phase a <b>liquid</b> (on an (inert) solid) / a solid (1)</p> <p>HPLC: <b>stationary</b> phase a <b>solid</b> / silica (1)</p> <p>HPLC: <b>mobile</b> phase a <b>liquid</b> (1)</p>	<p>Different retention times without a reason why</p> <p>Different volatilities</p> <p>Different masses</p> <p>Different reactivity</p> <p>Different reactions</p> <p>Different interactions</p>	5

TOTAL FOR SECTION C = 20 MARKS

Further copies of this publication are available from  
Edexcel Publications, Adamsway, Mansfield, Notts, NG18 4FN

Telephone 01623 467467  
Fax 01623 450481

Email [publications@linneydirect.com](mailto:publications@linneydirect.com)

Order Code UA026200 January 2011

For more information on Edexcel qualifications, please visit [www.edexcel.com/quals](http://www.edexcel.com/quals)

Edexcel Limited. Registered in England and Wales no.4496750  
Registered Office: One90 High Holborn, London, WC1V 7BH